


AMENDMENT TO THE CLAIMS

A listing of the claims presented in this patent application appears below. This listing replaces all prior versions and listing of claims in this patent application.

Claims 1-40 (canceled).

Claim 41 (currently amended): A single compound of the formula:



wherein:

R¹, R², R³ and R⁴ are, independently, selected from the group consisting of a hydrogen atom, halo, hydroxy, protected hydroxy, cyano, C₁ to C₁₂ alkyl, C₂ to C₁₂ alkenyl, C₂ to C₁₂ alkynyl, C₁ to C₁₂ substituted alkyl, C₂ to C₁₂ substituted alkenyl, C₂ to C₁₂ substituted alkynyl, C₁ to C₁₂ alkoxy, C₁ to C₁₂ substituted alkoxy, C₁ to C₁₂ acyloxy, C₁ to C₁₂ acyl, C₃ to C₇ cycloalkyl, C₃ to C₇ substituted cycloalkyl, C₅ to C₇ cycloalkenyl, C₅ to C₇ substituted cycloalkenyl, heterocyclic ring, substituted heterocyclic ring, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, cyclic C₂ to C₇ alkylene, substituted cyclic C₂ to C₇ alkylene, cyclic C₂ to C₇ heteroalkylene, substituted cyclic C₂ to C₇

heteroalkylene, carboxy, protected carboxy, hydroxymethyl, protected hydroxymethyl, protected amino, (monosubstituted)amino, protected (monosubstituted)amino, (disubstituted)amino, C₁ to C₁₀ alkylamino, C₁ to C₁₀ substituted alkylamino, carboxamide, protected carboxamide, C₁ to C₁₀ alkylthio, C₁ to C₁₀ substituted alkylthio, C₁ to C₁₀ alkylsulfonyl, C₁ to C₁₀ substituted alkylsulfonyl, C₁ to C₁₀ alkylsulfoxide, C₁ to C₁₀ substituted alkylsulfoxide, phenylthio, substituted phenylthio, phenylsulfoxide, substituted phenylsulfoxide, phenylsulfonyl, substituted phenylsulfonyl and the group consisting of (i) the formula $-C(O)NR^{11}R^{12}$, (ii) the formula $-C(O)R^{11}$, (iii) the formula $-NR^{11}R^{12}$, (iv) the formula $-SR^{11}$, (v) the formula $-OR^{11}$ and (vi) the formula $-C(O)OR^{11}$, wherein R¹¹ and R¹² are, independently, selected from the group consisting of a hydrogen atom, C₁ to C₁₂ alkyl, C₁ to C₁₂ substituted alkyl, C₂ to C₁₂ alkenyl, C₂ to C₁₂ substituted alkenyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, heterocycle, substituted heterocycle, phenylsulfonyl, substituted phenylsulfonyl, C₁ to C₁₀ alkylsulfonyl, C₁ to C₁₀ substituted alkylsulfonyl, C₁ to C₁₂ alkylaminocarbonyl, C₁ to C₁₂ substituted alkylaminocarbonyl, phenylaminocarbonyl and substituted phenylaminocarbonyl;

R⁵ is selected from the group consisting of a hydrogen atom, C₁ to C₁₂ alkyl, C₁ to C₁₂ substituted alkyl, phenyl, substituted phenyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, carboxy, protected carboxy, cyano, protected (monosubstituted)amino, (disubstituted)amino, C₁ to C₁₂ acyl, C₁ to C₁₂ substituted acyl, C₁ to C₁₂ alkoxycarbonyl, C₁ to C₁₂ substituted alkoxycarbonyl, heterocycle, substituted heterocycle, naphthyl, substituted naphthyl, C₃ to C₇ cycloalkyl, C₃ to C₇ substituted cycloalkyl, C₅ to C₇ cycloalkenyl and C₅ to C₇ substituted cycloalkenyl;


R⁶ is the formula:

-D-W-E-

wherein:

W is selected from the group consisting of phenylene, substituted phenylene, C₃ to C₇ cycloalkylene, C₃ to C₇ substituted cycloalkylene, C₅ to C₇ cycloalkenylene, C₅ to C₇ substituted cycloalkenylene, arylene, substituted arylene, heterocyclene, substituted heterocyclene, heteroarylene and substituted heteroarylene; and

D, which is directly attached to the nitrogen depicted in the formula, and E, which can be absent, are independently selected from the group consisting of C₁ to C₁₂ alkylene, C₂ to C₁₂ alkenylene, C₂ to C₁₂ alkynylene, C₁ to C₁₂ substituted alkylene, C₂ to C₁₂ substituted alkenylene, C₂ to C₁₂ substituted alkynylene, C₃ to C₇ cycloalkylene, C₃ to C₇ substituted cycloalkylene, C₅ to C₇ cycloalkenylene, C₅ to C₇ substituted cycloalkenylene, C₇ to C₁₈ phenylalkylene, C₇ to C₁₈ substituted phenylalkylene, C₁ to C₁₂ heterocycloalkylene and C₁ to C₁₂ substituted heterocycloalkylene, -NH- and the formula:



wherein R⁹ and R¹⁰ are, independently, selected from the group consisting of a hydrogen atom, C₁ to C₁₂ alkyl, C₂ to C₁₂ alkenyl, C₂ to C₁₂ alkynyl, C₁ to C₁₂ substituted alkyl, C₂ to C₁₂ substituted alkenyl, C₂ to C₁₂ substituted alkynyl, C₁ to C₁₂ acyl, C₁ to C₁₂ substituted acyl, C₃ to C₇ cycloalkyl, C₃ to C₇ substituted cycloalkyl, C₅ to C₇ cycloalkenyl, C₅ to C₇ substituted cycloalkenyl, a heterocyclic ring, substituted heterocyclic ring, heteroaryl, substituted heteroaryl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, C₇ to C₁₈ phenylalkoxy, C₇ to C₁₈ substituted phenylalkoxy, phenyl, substituted phenyl, naphthyl, substituted naphthyl, cyclic C₂ to C₇ alkylene, substituted cyclic C₂ to C₇

alkylene, cyclic C₂ to C₇ heteroalkylene, substituted cyclic C₂ to C₇ heteroalkylene, carboxy, protected carboxy, hydroxymethyl and protected hydroxymethyl; and m and n are, independently, 0, 1, 2, 3 or 4; and

R⁷ and R⁸ are, independently, selected from the group consisting of a functionalized resin, a hydrogen atom, C₁ to C₁₂ alkyl, C₁ to C₁₂ substituted alkyl, phenyl, substituted phenyl, heterocycle, substituted heterocycle, C₃ to C₇ cycloalkyl, C₃ to C₇ substituted cycloalkyl, C₅ to C₇ cycloalkenyl, C₅ to C₇ substituted cycloalkenyl, C₂ to C₁₂ alkenyl, C₂ to C₁₂ substituted alkenyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, [[and]] C₁ to C₁₂ substituted heterocycloalkyl, C₁ to C₁₂ acyl, C₁ to C₁₂ substituted acyl, phenylsulfonyl, substituted phenylsulfonyl, C₁ to C₁₀ alkylsulfonyl, C₁ to C₁₀ substituted alkylsulfonyl, C₁ to C₁₂ alkylaminocarbonyl, C₁ to C₁₂ substituted alkylaminocarbonyl, phenylaminocarbonyl, substituted phenylaminocarbonyl, C₁ to C₁₂ alkylaminothiocarbonyl, C₁ to C₁₂ substituted alkylaminothiocarbonyl, phenylaminothiocarbonyl and substituted phenylaminothiocarbonyl;


provided that, where R⁶ is methylene, at least one of R¹ to R⁴ must be the formula -C(O)NR¹¹R¹²; or

provided that, where R⁶ is methylene, at least one of R¹ to R⁴ must be the formula -C(O)R¹¹, wherein R¹¹ is a heterocyclic ring or substituted heterocyclic ring, wherein said ring contains at least one nitrogen atom and wherein said nitrogen atom is attached to the carbonyl carbon; or

a pharmaceutically acceptable salt of a compound thereof;

with the proviso that when R⁷ and R⁸ are hydrogen or -CH₂CH₃, substituents R¹, R², R³ and R⁴ cannot be hydrogen.

Claim 42 (currently amended): A single compound of the formula:



wherein:

R¹, R² and R⁴ are, independently, selected from the group consisting of a hydrogen atom, halo, hydroxy, protected hydroxy, cyano, C₁ to C₁₂ alkyl, C₂ to C₁₂ alkenyl, C₂ to C₁₂ alkynyl, C₁ to C₁₂ substituted alkyl, C₂ to C₁₂ substituted alkenyl, C₂ to C₁₂ substituted alkynyl, C₁ to C₁₂ alkoxy, C₁ to C₁₂ substituted alkoxy, C₁ to C₁₂ acyloxy, C₁ to C₁₂ acyl, C₃ to C₇ cycloalkyl, C₃ to C₇ substituted cycloalkyl, C₅ to C₇ cycloalkenyl, C₅ to C₇ substituted cycloalkenyl, heterocyclic ring, substituted heterocyclic ring, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, cyclic C₂ to C₇ alkylene, substituted cyclic C₂ to C₇ alkylene, cyclic C₂ to C₇ heteroalkylene, substituted cyclic C₂ to C₇ heteroalkylene, carboxy, protected carboxy, hydroxymethyl, protected hydroxymethyl, protected amino, (monosubstituted)amino, protected (monosubstituted)amino, (disubstituted)amino, C₁ to C₁₀ alkylamino, C₁ to C₁₀ substituted alkylamino, carboxamide, protected carboxamide, C₁ to C₁₀ alkylthio, C₁ to C₁₀ substituted alkylthio, C₁ to C₁₀ alkylsulfonyl, C₁ to C₁₀ substituted alkylsulfonyl, C₁ to C₁₀ alkylsulfoxide, C₁ to C₁₀ substituted alkylsulfoxide, phenylthio, substituted phenylthio, phenylsulfoxide, substituted phenylsulfoxide, phenylsulfonyl, substituted phenylsulfonyl and the group consisting of (i) the formula $-C(O)NR^{11}R^{12}$, (ii) the formula $-C(O)R^{11}$, (iii) the formula

$-NR^{11}R^{12}$, (iv) the formula $-SR^{11}$, (v) the formula $-OR^{11}$ and (vi) the formula $-C(O)OR^{11}$, wherein R^{11} and R^{12} are, independently, selected from the group consisting of a hydrogen atom, C_1 to C_{12} alkyl, C_1 to C_{12} substituted alkyl, C_2 to C_{12} alkenyl, C_2 to C_{12} substituted alkenyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, C_7 to C_{18} phenylalkyl, C_7 to C_{18} substituted phenylalkyl, C_1 to C_{12} heterocycloalkyl, C_1 to C_{12} substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, heterocycle, substituted heterocycle, phenylsulfonyl, substituted phenylsulfonyl, C_1 to C_{10} alkylsulfonyl, C_1 to C_{10} substituted alkylsulfonyl, C_1 to C_{12} alkylaminocarbonyl, C_1 to C_{12} substituted alkylaminocarbonyl, phenylaminocarbonyl and substituted phenylaminocarbonyl;

R^3 is selected from the group consisting of hydroxy, protected hydroxy, cyano, C_2 to C_{12} alkenyl, C_2 to C_{12} alkynyl, C_1 to C_{12} substituted alkyl, C_2 to C_{12} substituted alkenyl, C_2 to C_{12} substituted alkynyl, C_1 to C_{12} alkoxy, C_1 to C_{12} substituted alkoxy, C_1 to C_{12} acyloxy, C_1 to C_{12} acyl, C_3 to C_7 cycloalkyl, C_3 to C_7 substituted cycloalkyl, C_5 to C_7 cycloalkenyl, C_5 to C_7 substituted cycloalkenyl, heterocyclic ring, substituted heterocyclic ring, C_7 to C_{18} phenylalkyl, C_7 to C_{18} substituted phenylalkyl, C_1 to C_{12} heterocycloalkyl, C_1 to C_{12} substituted heterocycloalkyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, cyclic C_2 to C_7 alkylene, substituted cyclic C_2 to C_7 alkylene, cyclic C_2 to C_7 heteroalkylene, substituted cyclic C_2 to C_7 heteroalkylene, carboxy, protected carboxy, hydroxymethyl, protected hydroxymethyl, protected amino, (monosubstituted)amino, protected (monosubstituted)amino, (disubstituted)amino, C_1 to C_{10} alkylamino, C_1 to C_{10} substituted alkylamino, carboxamide, protected carboxamide, C_1 to C_{10} alkylthio, C_1 to C_{10} substituted alkylthio, C_1 to C_{10} alkylsulfonyl, C_1 to C_{10} substituted alkylsulfonyl, C_1 to C_{10} alkylsulfoxide, C_1 to C_{10} substituted alkylsulfoxide, phenylthio, substituted phenylthio, phenylsulfoxide, substituted phenylsulfoxide, phenylsulfonyl, substituted phenylsulfonyl and the group consisting of (i) the formula $-C(O)NR^{11}R^{12}$, (ii) the formula $-C(O)R^{11}$, (iii) the formula $-NR^{11}R^{12}$, (iv) the formula $-SR^{11}$, (v) the formula $-OR^{11}$ and (vi) the formula $-C(O)OR^{11}$, wherein R^{11} and R^{12} are, independently, selected from the group consisting of a hydrogen atom, C_1 to C_{12} alkyl, C_1 to C_{12} substituted alkyl, C_2 to C_{12} alkenyl, C_2 to C_{12} substituted alkenyl, phenyl, substituted phenyl, naphthyl,

substituted naphthyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, heterocycle, substituted heterocycle, phenylsulfonyl, substituted phenylsulfonyl, C₁ to C₁₀ alkylsulfonyl, C₁ to C₁₀ substituted alkylsulfonyl, C₁ to C₁₂ alkylaminocarbonyl, C₁ to C₁₂ substituted alkylaminocarbonyl, phenylaminocarbonyl and substituted phenylaminocarbonyl;

R⁵ is selected from the group consisting of a hydrogen atom, C₁ to C₁₂ alkyl, C₁ to C₁₂ substituted alkyl, phenyl, substituted phenyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, carboxy, protected carboxy, cyano, protected (monosubstituted)amino, (disubstituted)amino, C₁ to C₁₂ acyl, C₁ to C₁₂ substituted acyl, C₁ to C₁₂ alkoxycarbonyl, C₁ to C₁₂ substituted alkoxycarbonyl, heterocycle, substituted heterocycle, naphthyl, substituted naphthyl, C₃ to C₇ cycloalkyl, C₃ to C₇ substituted cycloalkyl, C₅ to C₇ cycloalkenyl and C₅ to C₇ substituted cycloalkenyl;

R⁶ is the formula:




wherein:

zero, one or two of D, W and E can be absent;

W, if present, is selected from the group consisting of phenylene, substituted phenylene, C₃ to C₇ cycloalkylene, C₃ to C₇ substituted cycloalkylene, C₅ to C₇ cycloalkenylene, C₅ to C₇ substituted cycloalkenylene, arylene, substituted arylene, heterocyclene, substituted heterocyclene, heteroarylene and substituted heteroarylene; and

D, which is directly attached to the nitrogen depicted in the formula, if present and E, if present, are independently selected from the group consisting of C₁ to C₁₂ alkylene, C₂ to C₁₂ alkenylene, C₂ to C₁₂ alkynylene, C₁ to C₁₂ substituted alkylene, C₂ to C₁₂ substituted alkenylene, C₂ to C₁₂ substituted alkynylene, C₃ to C₇ cycloalkylene, C₃ to C₇ substituted cycloalkylene, C₅ to C₇ cycloalkenylene, C₅ to C₇ substituted cycloalkenylene, C₇ to C₁₈ phenylalkylene, C₇ to C₁₈ substituted phenylalkylene, C₁ to C₁₂ heterocycloalkylene and C₁ to C₁₂ substituted heterocycloalkylene, -NH- and the formula:



wherein R^9 and R^{10} are, independently, selected from the group consisting of a hydrogen atom, C_1 to C_{12} alkyl, C_2 to C_{12} alkenyl, C_2 to C_{12} alkynyl, C_1 to C_{12} substituted alkyl, C_2 to C_{12} substituted alkenyl, C_2 to C_{12} substituted alkynyl, C_1 to C_{12} acyl, C_1 to C_{12} substituted acyl, C_3 to C_7 cycloalkyl, C_3 to C_7 substituted cycloalkyl, C_5 to C_7 cycloalkenyl, C_5 to C_7 substituted cycloalkenyl, a heterocyclic ring, substituted heterocyclic ring, heteroaryl, substituted heteroaryl, C_7 to C_{18} phenylalkyl, C_7 to C_{18} substituted phenylalkyl, C_1 to C_{12} heterocycloalkyl, C_1 to C_{12} substituted heterocycloalkyl, C_7 to C_{18} phenylalkoxy, C_7 to C_{18} substituted phenylalkoxy, phenyl, substituted phenyl, naphthyl, substituted naphthyl, cyclic C_2 to C_7 alkylene, substituted cyclic C_2 to C_7 alkylene, cyclic C_2 to C_7 heteroalkylene, substituted cyclic C_2 to C_7 heteroalkylene, carboxy, protected carboxy, hydroxymethyl and protected hydroxymethyl; and m and n are, independently, 0, 1, 2, 3 or 4; and

R^7 and R^8 are, independently, selected from the group consisting of a functionalized resin, a hydrogen atom, C_1 to C_{12} alkyl, C_1 to C_{12} substituted alkyl, phenyl, substituted phenyl, heterocycle, substituted heterocycle, C_3 to C_7 cycloalkyl, C_3 to C_7 substituted cycloalkyl, C_5 to C_7 cycloalkenyl, C_5 to C_7 substituted cycloalkenyl, C_2 to C_{12} alkenyl, C_2 to C_{12} substituted alkenyl, C_7 to C_{18} phenylalkyl, C_7 to C_{18} substituted phenylalkyl, C_1 to C_{12} heterocycloalkyl, [[and]] C_1 to C_{12} substituted heterocycloalkyl, C_1 to C_{12} acyl, C_1 to C_{12} substituted acyl, phenylsulfonyl, substituted phenylsulfonyl, C_1 to C_{10} alkylsulfonyl, C_1 to C_{10} substituted alkylsulfonyl, C_1 to C_{12} alkylaminocarbonyl, C_1 to C_{12} substituted alkylaminocarbonyl, phenylaminocarbonyl, substituted phenylaminocarbonyl, C_1 to C_{12}


alkylaminothiocarbonyl, C₁ to C₁₂ substituted alkylaminothiocarbonyl, phenylaminothiocarbonyl and substituted phenylaminothiocarbonyl; or

a pharmaceutically acceptable salt of a compound thereof;

with the proviso that when R⁷ and R⁸ are hydrogen or -CH₂CH₃, substituents R¹, R², R³ and R⁴ cannot be hydrogen.

Claim 43 (canceled).

Claim 44 (currently amended): A single compound of the formula:



wherein:

R¹, R², R³ and R⁴ are, independently, selected from the group consisting of a hydrogen atom, halo, hydroxy, protected hydroxy, cyano, C₁ to C₁₂ alkyl, C₂ to C₁₂ alkenyl, C₂ to C₁₂ alkynyl, C₁ to C₁₂ substituted alkyl, C₂ to C₁₂ substituted alkenyl, C₂ to C₁₂ substituted alkynyl, C₁ to C₁₂ alkoxy, C₁ to C₁₂ substituted alkoxy, C₁ to C₁₂ acyloxy, C₁ to C₁₂ acyl, C₃ to C₇ cycloalkyl, C₃ to C₇ substituted cycloalkyl, C₅ to C₇ cycloalkenyl, C₅ to C₇ substituted cycloalkenyl, heterocyclic ring, substituted heterocyclic ring, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, cyclic C₂ to C₇ alkylene, substituted

cyclic C₂ to C₇ alkylene, cyclic C₂ to C₇ heteroalkylene, substituted cyclic C₂ to C₇ heteroalkylene, carboxy, protected carboxy, hydroxymethyl, protected hydroxymethyl, protected amino, (monosubstituted)amino, protected (monosubstituted)amino, (disubstituted)amino, C₁ to C₁₀ alkylamino, C₁ to C₁₀ substituted alkylamino, carboxamide, protected carboxamide, C₁ to C₁₀ alkylthio, C₁ to C₁₀ substituted alkylthio, C₁ to C₁₀ alkylsulfonyl, C₁ to C₁₀ substituted alkylsulfonyl, C₁ to C₁₀ alkylsulfoxide, C₁ to C₁₀ substituted alkylsulfoxide, phenylthio, substituted phenylthio, phenylsulfoxide, substituted phenylsulfoxide, phenylsulfonyl, substituted phenylsulfonyl and the group consisting of (i) the formula $-C(O)NR^{11}R^{12}$, (ii) the formula $-C(O)R^{11}$, (iii) the formula $-NR^{11}R^{12}$, (iv) the formula $-SR^{11}$, (v) the formula $-OR^{11}$ and (vi) the formula $-C(O)OR^{11}$, wherein R¹¹ and R¹² are, independently, selected from the group consisting of a hydrogen atom, C₁ to C₁₂ alkyl, C₁ to C₁₂ substituted alkyl, C₂ to C₁₂ alkenyl, C₂ to C₁₂ substituted alkenyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, heterocycle, substituted heterocycle, phenylsulfonyl, substituted phenylsulfonyl, C₁ to C₁₀ alkylsulfonyl, C₁ to C₁₀ substituted alkylsulfonyl, C₁ to C₁₂ alkylaminocarbonyl, C₁ to C₁₂ substituted alkylaminocarbonyl, phenylaminocarbonyl and substituted phenylaminocarbonyl;

R⁵ is selected from the group consisting of phenyl, substituted phenyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, carboxy, protected carboxy, protected (monosubstituted)amino, (disubstituted)amino, C₁ to C₁₂ substituted acyl, C₁ to C₁₂ alkoxycarbonyl, C₁ to C₁₂ substituted alkoxycarbonyl, heterocycle, substituted heterocycle, naphthyl, substituted naphthyl, C₃ to C₇ cycloalkyl, C₃ to C₇ substituted cycloalkyl, C₅ to C₇ cycloalkenyl and C₅ to C₇ substituted cycloalkenyl;

R⁶ is the formula:


-D-W-E-

wherein:

zero, one or two of D, W, and E can be absent;

W, if present, is selected from the group consisting of phenylene, substituted phenylene, C₃ to C₇ cycloalkylene, C₃ to C₇ substituted cycloalkylene, C₅ to C₇ cycloalkenylene, C₅ to C₇ substituted cycloalkenylene, arylene, substituted arylene, heterocyclene, substituted heterocyclene, heteroarylene and substituted heteroarylene; and

D, which is directly attached to the nitrogen depicted in the formula, if present, and E, if present, are independently selected from the group consisting of C₁ to C₁₂ alkylene, C₂ to C₁₂ alkenylene, C₂ to C₁₂ alkynylene, C₁ to C₁₂ substituted alkylene, C₂ to C₁₂ substituted alkenylene, C₂ to C₁₂ substituted alkynylene, C₃ to C₇ cycloalkylene, C₃ to C₇ substituted cycloalkylene, C₅ to C₇ cycloalkenylene, C₅ to C₇ substituted cycloalkenylene, C₇ to C₁₈ phenylalkylene, C₇ to C₁₈ substituted phenylalkylene, C₁ to C₁₂ heterocycloalkylene and C₁ to C₁₂ substituted heterocycloalkylene, -NH- and the formula:



wherein R⁹ and R¹⁰ are, independently, selected from the group consisting of a hydrogen atom, C₁ to C₁₂ alkyl, C₂ to C₁₂ alkenyl, C₂ to C₁₂ alkynyl, C₁ to C₁₂ substituted alkyl, C₂ to C₁₂ substituted alkenyl, C₂ to C₁₂ substituted alkynyl, C₁ to C₁₂ acyl, C₁ to C₁₂ substituted acyl, C₃ to C₇ cycloalkyl, C₃ to C₇ substituted cycloalkyl, C₅ to C₇ cycloalkenyl, C₅ to C₇ substituted cycloalkenyl, a heterocyclic ring, substituted heterocyclic ring, heteroaryl, substituted heteroaryl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, C₇ to C₁₈ phenylalkoxy, C₇ to C₁₈ substituted phenylalkoxy, phenyl, substituted phenyl, naphthyl, substituted naphthyl, cyclic C₂ to C₇ alkylene, substituted cyclic C₂ to C₇

alkylene, cyclic C₂ to C₇ heteroalkylene, substituted cyclic C₂ to C₇ heteroalkylene, carboxy, protected carboxy, hydroxymethyl and protected hydroxymethyl; and m and n are, independently, 0, 1, 2, 3 or 4; and R⁷ and R⁸ are, independently, selected from the group consisting of a functionalized resin, a hydrogen atom, C₁ to C₁₂ alkyl, C₁ to C₁₂ substituted alkyl, phenyl, substituted phenyl, heterocycle, substituted heterocycle, C₃ to C₇ cycloalkyl, C₃ to C₇ substituted cycloalkyl, C₅ to C₇ cycloalkenyl, C₅ to C₇ substituted cycloalkenyl, C₂ to C₁₂ alkenyl, C₂ to C₁₂ substituted alkenyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, [[and]] C₁ to C₁₂ substituted heterocycloalkyl, C₁ to C₁₂ acyl, C₁ to C₁₂ substituted acyl, phenylsulfonyl, substituted phenylsulfonyl, C₁ to C₁₀ alkylsulfonyl, C₁ to C₁₀ substituted alkylsulfonyl, C₁ to C₁₂ alkylaminocarbonyl, C₁ to C₁₂ substituted alkylaminocarbonyl, phenylaminocarbonyl, substituted phenylaminocarbonyl, C₁ to C₁₂ alkylaminothiocarbonyl, C₁ to C₁₂ substituted alkylaminothiocarbonyl, phenylaminothiocarbonyl and substituted phenylaminothiocarbonyl; or
a pharmaceutically acceptable salt of a compound thereof;
with the proviso that when R⁷ and R⁸ are hydrogen or -CH₂CH₃, substituents R¹, R², R³ and R⁴ cannot be hydrogen.

Claim 45 (canceled).

Claim 46 (previously added): The single compound of claim 44, wherein:

R¹, R² and R⁴ are each a hydrogen atom and R³ is the formula -C(O)NR¹¹R¹², wherein R¹¹ is selected from the group consisting of a hydrogen atom, methyl, ethyl and benzyl and R¹² is selected from the group consisting of a hydrogen atom, 2-(2-methoxyphenyl)ethyl, (1-ethyl-2-pyrrolidino)methyl, pyridin-2-ylmethyl, 2-methyl-5-chlorophenyl, (2-(pyridin-2-yl)ethyl), 1-ethyl-2-pyrrolidinylmethyl, 3,3,5-trimethylcyclohexyl, 3,4-methylenedioxyphenyl, 3-(trifluoromethyl)benzyl, pyridin-4-ylmethyl, 6-indazolyl, 2-(ethoxycarbonyl)ethyl, cyclooctyl, cyclopropyl, benzyl, N,N-(diethylamino)ethyl,

3-(2-oxo-1-pyrrolidine)propyl, 3-(4-morpholino)propyl, (ethoxycarbonyl)methyl and cyclohexyl;

R⁵ is selected from the group consisting of phenoxyphenyl, 4-hydroxy-3-methoxyphenyl, 3,4,5-trimethoxyphenyl, 3-hydroxy-4-methoxyphenyl, 4-acetamidophenyl, 4-phenoxyphenyl, 4-methoxyl-1-naphthyl, 4-bromo-2-thienyl, 4-pyridyl, isopropyl, 2-methylthioethyl, 4-chloro-3-nitrophenyl, 3-nitrophenyl, 4-t-butylphenyl, 2,3-dichlorophenyl, 3,5-bis(trifluoromethyl)phenyl, 2,5-difluorophenyl, 2-quinolyl, 2-chloro-3,4-dimethoxyphenyl, 5-methyl-2-furyl, 4-chloro-3-fluorophenyl, 2-phenyl-4-imidazolyl, 2-(ethoxycarbonyl)cyclopropyl, 5-nitro-2-furyl, 4-bromophenyl, cyclopropyl, 2-norbornen-5-yl, 6-nitropiperonyl, 2-chloro-5-nitrophenyl, 5-hydroxy-2-nitrophenyl, 3-hydroxyphenyl, 3,4-difluorophenyl, 4-dimethylaminophenyl, 4-methylthiophenyl, 4-(trifluoromethyl)phenyl, 2-thienyl, 2,3-dimethoxyphenyl, 3-ethoxy-4-hydroxyphenyl, 4-cyanophenyl, 3-cyanophenyl, 2-furyl, 4-nitrophenyl, 1-naphthyl, 2-methoxyphenyl, 4-isopropylphenyl, piperonyl, 2-fluorophenyl, 4-ethoxyphenyl and 2,4-dihydroxyphenyl;

R⁶ is selected from the group consisting of methylene, ethylidene, ethylene, propylene, pentylene, isopentylidene, 3-aminocarbonylbutylidene, 2-methylthiopropylidene, isobutylidene, phenylmethylene, benzylmethylene, cyclohexylethylidene, 4-chlorobenzylmethylene, indol-3-ylethylidene, 4-trifluoroacetamidopentylidene, 3-guanidobutylidene, hydroxyethylidene, 2-aminocarbonylpropylidene, isopentylidene, mercaptoethylidene, 4-hydroxybenzylmethylene, 1,3-phenylene, 1,4-phenylene, 1,4-(phenylene)-NH-, 3,6-dioxaoctylene-NH-, -CH₂CH₂NH- and 1,4-(cyclohexylene)-NH-; and

R⁷ and R⁸ are each a hydrogen atom.

Claim 47 (previously added): The single compound of claim 44, wherein:

R¹, R² and R⁴ are each a hydrogen atom and R³ is the formula -C(O)R¹¹, wherein R¹¹ is selected from the group consisting of 1,3,3-trimethyl-6-aza-6-bicyclo(3,2,1)octyl, 4-(4-fluorophenyl)-1-piperazino, 4-acetyl-1-piperazino, piperazino,

2-methyl-4-(3-methylphenyl)-1-piperazino, 4-(ethoxycarbonyl)piperidino, N-methylhomopiperazino and N,N'-diisopropylimidamino;


R⁵ is selected from the group consisting of phenoxyphenyl, 4-hydroxy-3-methoxyphenyl, 3,4,5-trimethoxyphenyl, 3-hydroxy-4-methoxyphenyl, 4-acetamidophenyl, 4-phenoxyphenyl, 4-methoxyl-1-naphthyl, 4-bromo-2-thienyl, 4-pyridyl, isopropyl, 2-methylthioethyl, 4-chloro-3-nitrophenyl, 3-nitrophenyl, 4-t-butylphenyl, 2,3-dichlorophenyl, 3,5-bis(trifluoromethyl)phenyl, 2,5-difluorophenyl, 2-quinolyl, 2-chloro-3,4-dimethoxylphenyl, 5-methyl-2-furyl, 4-chloro-3-fluorophenyl, 2-phenyl-4-imidazolyl, 2-(ethoxycarbonyl)cyclopropyl, 5-nitro-2-furyl, 4-bromophenyl, cyclopropyl, 2-norbornen-5-yl, 6-nitropiperonyl, 2-chloro-5-nitrophenyl, 5-hydroxy-2-nitrophenyl, 3-hydroxyphenyl, 3,4-difluorophenyl, 4-dimethylaminophenyl, 4-methylthiophenyl, 4-(trifluoromethyl)phenyl, 2-thienyl, 2,3-dimethoxyphenyl, 3-ethoxy-4-hydroxyphenyl, 4-cyanophenyl, 3-cyanophenyl, 2-furyl, 4-nitrophenyl, 1-naphthyl, 2-methoxyphenyl, 4-isopropylphenyl, piperonyl, 2-fluorophenyl, 4-ethoxyphenyl and 2,4-dihydroxyphenyl;

R⁶ is selected from the group consisting of methylene, ethylidene, ethylene, propylene, pentylene, isopentylidene, 3-aminocarbonylbutylidene, 2-methylthiopropylidene, isobutylidene, phenylmethylene, benzylmethylene, cyclohexylethylidene, 4-chlorobenzylmethylene, indol-3-ylethylidene, 4-trifluoroacetamidopentylidene, 3-guanidobutylidene, hydroxyethylidene, 2-aminocarbonylpropylidene, isopentylidene, mercaptoethylidene, 4-hydroxybenzylmethylene, 1,3-phenylene, 1,4-phenylene, 1,4-(phenylene)-NH-, 3,6-dioxaoctylene-NH-, -CH₂CH₂NH- and 1,4-(cyclohexylene)-NH-; and

R⁷ and R⁸ are each a hydrogen atom.

Claim 48 (canceled).

Claim 49 (new): A single compound of the formula:



wherein:

R¹, R² and R⁴ are, independently, selected from the group consisting of a hydrogen atom, halo, hydroxy, protected hydroxy, cyano, C₁ to C₁₂ alkyl, C₂ to C₁₂ alkenyl, C₂ to C₁₂ alkynyl, C₁ to C₁₂ substituted alkyl, C₂ to C₁₂ substituted alkenyl, C₂ to C₁₂ substituted alkynyl, C₁ to C₁₂ alkoxy, C₁ to C₁₂ substituted alkoxy, C₁ to C₁₂ acyloxy, C₁ to C₁₂ acyl, C₃ to C₇ cycloalkyl, C₃ to C₇ substituted cycloalkyl, C₅ to C₇ cycloalkenyl, C₅ to C₇ substituted cycloalkenyl, heterocyclic ring, substituted heterocyclic ring, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, cyclic C₂ to C₇ alkylene, substituted cyclic C₂ to C₇ alkylene, cyclic C₂ to C₇ heteroalkylene, substituted cyclic C₂ to C₇ heteroalkylene, carboxy, protected carboxy, hydroxymethyl, protected hydroxymethyl, protected amino, (monosubstituted)amino, protected (monosubstituted)amino, (disubstituted)amino, C₁ to C₁₀ alkylamino, C₁ to C₁₀ substituted alkylamino, carboxamide, protected carboxamide, C₁ to C₁₀ alkylthio, C₁ to C₁₀ substituted alkylthio, C₁ to C₁₀ alkylsulfonyl, C₁ to C₁₀ substituted alkylsulfonyl, C₁ to C₁₀ alkylsulfoxide, C₁ to C₁₀ substituted alkylsulfoxide, phenylthio, substituted phenylthio, phenylsulfoxide, substituted phenylsulfoxide, phenylsulfonyl, substituted phenylsulfonyl and the

group consisting of (i) the formula $-C(O)NR^{11}R^{12}$, (ii) the formula $-C(O)R^{11}$, (iii) the formula $-NR^{11}R^{12}$, (iv) the formula $-SR^{11}$, (v) the formula $-OR^{11}$ and (vi) the formula $-C(O)OR^{11}$, wherein R^{11} and R^{12} are, independently, selected from the group consisting of a hydrogen atom, C_1 to C_{12} alkyl, C_1 to C_{12} substituted alkyl, C_2 to C_{12} alkenyl, C_2 to C_{12} substituted alkenyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, C_7 to C_{18} phenylalkyl, C_7 to C_{18} substituted phenylalkyl, C_1 to C_{12} heterocycloalkyl, C_1 to C_{12} substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, heterocycle, substituted heterocycle, phenylsulfonyl, substituted phenylsulfonyl, C_1 to C_{10} alkylsulfonyl, C_1 to C_{10} substituted alkylsulfonyl, C_1 to C_{12} alkylaminocarbonyl, C_1 to C_{12} substituted alkylaminocarbonyl, phenylaminocarbonyl and substituted phenylaminocarbonyl;

R^3 is selected from the group consisting of hydroxy, protected hydroxy, cyano, C_2 to C_{12} alkenyl, C_2 to C_{12} alkynyl, C_1 to C_{12} substituted alkyl, C_2 to C_{12} substituted alkenyl, C_2 to C_{12} substituted alkynyl, C_1 to C_{12} alkoxy, C_1 to C_{12} substituted alkoxy, C_1 to C_{12} acyloxy, C_1 to C_{12} acyl, C_3 to C_7 cycloalkyl, C_3 to C_7 substituted cycloalkyl, C_5 to C_7 cycloalkenyl, C_5 to C_7 substituted cycloalkenyl, heterocyclic ring, substituted heterocyclic ring, C_7 to C_{18} phenylalkyl, C_7 to C_{18} substituted phenylalkyl, C_1 to C_{12} heterocycloalkyl, C_1 to C_{12} substituted heterocycloalkyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, cyclic C_2 to C_7 alkylene, substituted cyclic C_2 to C_7 alkylene, cyclic C_2 to C_7 heteroalkylene, substituted cyclic C_2 to C_7 heteroalkylene, carboxy, protected carboxy, hydroxymethyl, protected hydroxymethyl, protected amino, (monosubstituted)amino, protected (monosubstituted)amino, (disubstituted)amino, C_1 to C_{10} alkylamino, C_1 to C_{10} substituted alkylamino, carboxamide, protected carboxamide, C_1 to C_{10} alkylthio, C_1 to C_{10} substituted alkylthio, C_1 to C_{10} alkylsulfonyl, C_1 to C_{10} substituted alkylsulfonyl, C_1 to C_{10} alkylsulfoxide, C_1 to C_{10} substituted alkylsulfoxide, phenylthio, substituted phenylthio, phenylsulfoxide, substituted phenylsulfoxide, phenylsulfonyl, substituted phenylsulfonyl and the group consisting of (i) the formula $-C(O)NR^{11}R^{12}$, (ii) the formula $-C(O)R^{11}$, (iii) the formula $-NR^{11}R^{12}$, (iv) the formula $-SR^{11}$, (v) the formula $-OR^{11}$ and (vi) the formula $-C(O)OR^{11}$, wherein R^{11} and R^{12} are, independently, selected from the group consisting of a hydrogen atom, C_1 to C_{12} alkyl, C_1 to C_{12} substituted

alkyl, C₂ to C₁₂ alkenyl, C₂ to C₁₂ substituted alkenyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, heterocycle, substituted heterocycle, phenylsulfonyl, substituted phenylsulfonyl, C₁ to C₁₀ alkylsulfonyl, C₁ to C₁₀ substituted alkylsulfonyl, C₁ to C₁₂ alkylaminocarbonyl, C₁ to C₁₂ substituted alkylaminocarbonyl, phenylaminocarbonyl and substituted phenylaminocarbonyl;

R⁵ is selected from the group consisting of a hydrogen atom, C₁ to C₁₂ alkyl, C₁ to C₁₂ substituted alkyl, phenyl, substituted phenyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, carboxy, protected carboxy, cyano, protected (monosubstituted)amino, (disubstituted)amino, C₁ to C₁₂ acyl, C₁ to C₁₂ substituted acyl, C₁ to C₁₂ alkoxy carbonyl, C₁ to C₁₂ substituted alkoxy carbonyl, heterocycle, substituted heterocycle, naphthyl, substituted naphthyl, C₃ to C₇ cycloalkyl, C₃ to C₇ substituted cycloalkyl, C₅ to C₇ cycloalkenyl and C₅ to C₇ substituted cycloalkenyl;


R⁶ is the formula:

-D-W-E-

wherein:

W is absent or selected from the group consisting of phenylene, substituted phenylene, C₃ to C₇ cycloalkylene, C₃ to C₇ substituted cycloalkylene, C₅ to C₇ cycloalkenylene, C₅ to C₇ substituted cycloalkenylene, arylene, substituted arylene, heterocyclene, substituted heterocyclene, heteroarylene and substituted heteroarylene; and

D, which is directly attached to the nitrogen depicted in the formula, and E, which can be absent, are independently selected from the group consisting of C₁ to C₁₂ alkylene, C₂ to C₁₂ alkenylene, C₂ to C₁₂ alkynylene, C₁ to C₁₂ substituted alkylene, C₂ to C₁₂ substituted alkenylene, C₂ to C₁₂ substituted alkynylene, C₃ to C₇ cycloalkylene, C₃ to C₇ substituted cycloalkylene, C₅ to C₇ cycloalkenylene, C₅ to C₇ substituted cycloalkenylene, C₇ to C₁₈ phenylalkylene, C₇ to C₁₈ substituted phenylalkylene, C₁ to C₁₂ heterocycloalkylene and C₁ to C₁₂ substituted heterocycloalkylene, -NH- and the formula:



wherein R^9 and R^{10} are, independently, selected from the group consisting of a hydrogen atom, C_1 to C_{12} alkyl, C_2 to C_{12} alkenyl, C_2 to C_{12} alkynyl, C_1 to C_{12} substituted alkyl, C_2 to C_{12} substituted alkenyl, C_2 to C_{12} substituted alkynyl, C_1 to C_{12} acyl, C_1 to C_{12} substituted acyl, C_3 to C_7 cycloalkyl, C_3 to C_7 substituted cycloalkyl, C_5 to C_7 cycloalkenyl, C_5 to C_7 substituted cycloalkenyl, a heterocyclic ring, substituted heterocyclic ring, heteroaryl, substituted heteroaryl, C_7 to C_{18} phenylalkyl, C_7 to C_{18} substituted phenylalkyl, C_1 to C_{12} heterocycloalkyl, C_1 to C_{12} substituted heterocycloalkyl, C_7 to C_{18} phenylalkoxy, C_7 to C_{18} substituted phenylalkoxy, phenyl, substituted phenyl, naphthyl, substituted naphthyl, cyclic C_2 to C_7 alkylene, substituted cyclic C_2 to C_7 alkylene, cyclic C_2 to C_7 heteroalkylene, substituted cyclic C_2 to C_7 heteroalkylene, carboxy, protected carboxy, hydroxymethyl and protected hydroxymethyl; and m and n are, independently, 0, 1, 2, 3 or 4; and

R^7 and R^8 are, independently, selected from the group consisting of a functionalized resin, a hydrogen atom, C_1 to C_{12} alkyl, C_1 to C_{12} substituted alkyl, phenyl, substituted phenyl, heterocycle, substituted heterocycle, C_3 to C_7 cycloalkyl, C_3 to C_7 substituted cycloalkyl, C_5 to C_7 cycloalkenyl, C_5 to C_7 substituted cycloalkenyl, C_2 to C_{12} alkenyl, C_2 to C_{12} substituted alkenyl, C_7 to C_{18} phenylalkyl, C_7 to C_{18} substituted phenylalkyl, C_1 to C_{12} heterocycloalkyl, [[and]] C_1 to C_{12} substituted heterocycloalkyl, C_1 to C_{12} acyl, C_1 to C_{12} substituted acyl, phenylsulfonyl, substituted phenylsulfonyl, C_1 to C_{10} alkylsulfonyl, C_1 to C_{10} substituted alkylsulfonyl, C_1 to C_{12} alkylaminocarbonyl, C_1 to C_{12} substituted alkylaminocarbonyl, phenylaminocarbonyl, substituted phenylaminocarbonyl, C_1 to C_{12} alkylaminothiocarbonyl, C_1 to

C₁₂ substituted alkylaminothiocarbonyl, phenylaminothiocarbonyl and substituted phenylaminothiocarbonyl;

provided that, where R⁶ is methylene, at least one of R¹ to R⁴ must be the formula - C(O)NR¹¹R¹²; or

provided that, where R⁶ is methylene, at least one of R¹ to R⁴ must be the formula - C(O)R¹¹, wherein R¹¹ is a heterocyclic ring or substituted heterocyclic ring, wherein said ring contains at least one nitrogen atom and wherein said nitrogen atom is attached to the carbonyl carbon; or

a pharmaceutically acceptable salt of a compound thereof;

with the proviso that when R⁷ and R⁸ are hydrogen or -CH₂CH₃, substituents R¹, R², R³ and R⁴ cannot be hydrogen.

Claim 50 (new): The single compound of claim 49, wherein R⁵ is selected from the group consisting of a hydrogen atom, C₁ to C₁₂ alkyl, C₁ to C₁₂ substituted alkyl, phenyl, substituted phenyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, heterocycle, substituted heterocycle, C₃ to C₇ cycloalkyl and C₃ to C₇ substituted cycloalkyl.

Claim 51 (new): The single compound of claim 49, wherein:

R¹, R² and R³ are, independently, selected from the group consisting of a hydrogen atom, halo, C₁ to C₁₂ alkyl, C₁ to C₁₂ substituted alkyl, carboxy, and the group consisting of (i) the formula -C(O)NR¹¹R¹² and (ii) the formula -C(O)R¹¹, wherein R¹¹ and R¹² are, independently, selected from the group consisting of a hydrogen atom, C₁ to C₁₂ alkyl, C₁ to C₁₂ substituted alkyl, C₂ to C₁₂ alkenyl, C₂ to C₁₂ substituted alkenyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, heterocycle and substituted heterocycle;

R⁵ is selected from the group consisting of a hydrogen atom, C₁ to C₁₂ alkyl, C₁ to C₁₂ substituted alkyl, phenyl, substituted phenyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted

phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, heterocycle, substituted heterocycle, C₃ to C₇ cycloalkyl and C₃ to C₇ substituted cycloalkyl;


R⁶ is the formula:



wherein:

W is absent or selected from the group consisting of phenylene, substituted phenylene, C₃ to C₇ cycloalkylene and C₃ to C₇ substituted cycloalkylene; and

D, which is directly attached to the nitrogen depicted in the formula, and E, which can be absent, are, independently, selected from the group consisting of C₁ to C₁₂ alkylene, C₁ to C₁₂ substituted alkylene, -NH- and the formula:



wherein R⁹ and R¹⁰ are, independently, selected from the group consisting of a hydrogen atom, C₁ to C₁₂ alkyl, C₁ to C₁₂ substituted alkyl, C₃ to C₇ cycloalkyl, C₃ to C₇ substituted cycloalkyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, phenyl, substituted phenyl; and m and n are independently 0, 1 or 2; and

R⁷ and R⁸ are each a hydrogen atom.


Claim 52 (new): The single compound of claim 49, wherein R⁶ is methylene, R¹, R² and R⁴ are each a hydrogen atom and R³ is the formula -C(O)NR¹¹R¹².

Claim 53 (new): The single compound of claim 49, wherein R^6 is methylene, R^1 , R^2 and R^4 are each a hydrogen atom and R^3 is the formula $-C(O)R^{11}$, wherein R^{11} is a heterocyclic ring or substituted heterocyclic ring, wherein said ring contains at least one nitrogen atom and wherein said nitrogen atom is attached to the carbonyl carbon.

Claim 54 (new): The single compound of claim 49, wherein R^6 is not methylene.

Claim 55 (new): The single compound of claim 49, wherein R^3 is selected from (i) the formula $-C(O)NR^{11}R^{12}$ and (ii) the formula $-C(O)R^{11}$, wherein R^{11} and R^{12} are, independently, selected from the group consisting of a hydrogen atom, C_1 to C_{12} alkyl, C_1 to C_{12} substituted alkyl, C_2 to C_{12} alkenyl, C_2 to C_{12} substituted alkenyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, C_7 to C_{18} phenylalkyl, C_7 to C_{18} substituted phenylalkyl, C_1 to C_{12} heterocycloalkyl, C_1 to C_{12} substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, heterocycle, substituted heterocycle, phenylsulfonyl, substituted phenylsulfonyl, C_1 to C_{10} alkylsulfonyl, C_1 to C_{10} substituted alkylsulfonyl, C_1 to C_{12} alkylaminocarbonyl, C_1 to C_{12} substituted alkylaminocarbonyl, phenylaminocarbonyl and substituted phenylaminocarbonyl.

Claim 56 (new): A single compound of the formula:



wherein:

R^1 , R^2 , R^3 and R^4 are, independently, selected from the group consisting of a hydrogen atom, halo, hydroxy, protected hydroxy, cyano, C_1 to C_{12} alkyl, C_2 to C_{12} alkenyl, C_2 to C_{12} alkynyl, C_1 to C_{12} substituted alkyl, C_2 to C_{12} substituted alkenyl, C_2 to C_{12} substituted alkynyl, C_1 to C_{12} alkoxy, C_1 to C_{12} substituted alkoxy, C_1 to C_{12} acyloxy, C_1 to C_{12} acyl, C_3 to C_7 cycloalkyl, C_3 to C_7 substituted cycloalkyl, C_5 to C_7 cycloalkenyl, C_5 to C_7 substituted cycloalkenyl, heterocyclic ring, substituted heterocyclic ring, C_7 to C_{18} phenylalkyl, C_7 to C_{18} substituted phenylalkyl, C_1 to C_{12} heterocycloalkyl, C_1 to C_{12} substituted heterocycloalkyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, cyclic C_2 to C_7 alkylene, substituted cyclic C_2 to C_7 alkylene, cyclic C_2 to C_7 heteroalkylene, substituted cyclic C_2 to C_7 heteroalkylene, carboxy, protected carboxy, hydroxymethyl, protected hydroxymethyl, protected amino, (monosubstituted)amino, protected (monosubstituted)amino, (disubstituted)amino, C_1 to C_{10} alkylamino, C_1 to C_{10} substituted alkylamino, carboxamide, protected carboxamide, C_1 to C_{10} alkylthio, C_1 to C_{10} substituted alkylthio, C_1 to C_{10} alkylsulfonyl, C_1 to C_{10} substituted alkylsulfonyl, C_1 to C_{10} alkylsulfoxide, C_1 to C_{10} substituted alkylsulfoxide, phenylthio, substituted phenylthio, phenylsulfoxide, substituted phenylsulfoxide, phenylsulfonyl, substituted phenylsulfonyl and the group consisting of (i) the formula $-C(O)NR^{11}R^{12}$, (ii) the formula $-C(O)R^{11}$, (iii) the formula $-NR^{11}R^{12}$, (iv) the formula $-SR^{11}$, (v) the formula $-OR^{11}$ and (vi) the formula $-C(O)OR^{11}$, wherein R^{11} and R^{12} are, independently, selected from the group consisting of a hydrogen atom, C_1 to C_{12} alkyl, C_1 to C_{12} substituted alkyl, C_2 to C_{12} alkenyl, C_2 to C_{12} substituted alkenyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, C_7 to C_{18} phenylalkyl, C_7 to C_{18} substituted phenylalkyl, C_1 to C_{12} heterocycloalkyl, C_1 to C_{12} substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, heterocycle, substituted heterocycle, phenylsulfonyl, substituted phenylsulfonyl, C_1 to C_{10} alkylsulfonyl, C_1 to C_{10} substituted alkylsulfonyl, C_1 to C_{12} alkylaminocarbonyl, C_1 to C_{12} substituted alkylaminocarbonyl, phenylaminocarbonyl and substituted phenylaminocarbonyl;

R^5 is selected from the group consisting of phenyl, substituted phenyl, C_7 to C_{18} phenylalkyl, C_7 to C_{18} substituted phenylalkyl, C_1 to C_{12} heterocycloalkyl, C_1 to C_{12} substituted heterocycloalkyl, carboxy, protected carboxy, protected (monosubstituted)amino,

(disubstituted)amino, C₁ to C₁₂ alkoxycarbonyl, C₁ to C₁₂ substituted alkoxycarbonyl, heterocycle, substituted heterocycle, naphthyl, substituted naphthyl, C₃ to C₇ cycloalkyl, C₃ to C₇ substituted cycloalkyl, C₅ to C₇ cycloalkenyl and C₅ to C₇ substituted cycloalkenyl;


R⁶ is the formula:



wherein:

W is absent or selected from the group consisting of phenylene, substituted phenylene, C₃ to C₇ cycloalkylene, C₃ to C₇ substituted cycloalkylene, C₅ to C₇ cycloalkenylene, C₅ to C₇ substituted cycloalkenylene, arylene, substituted arylene, heterocyclene, substituted heterocyclene, heteroarylene and substituted heteroarylene; and

D, which is directly attached to the nitrogen depicted in the formula, and E, which can be absent, are independently selected from the group consisting of C₁ to C₁₂ alkylene, C₂ to C₁₂ alkenylene, C₂ to C₁₂ alkynylene, C₁ to C₁₂ substituted alkylene, C₂ to C₁₂ substituted alkenylene, C₂ to C₁₂ substituted alkynylene, C₃ to C₇ cycloalkylene, C₃ to C₇ substituted cycloalkylene, C₅ to C₇ cycloalkenylene, C₅ to C₇ substituted cycloalkenylene, C₇ to C₁₈ phenylalkylene, C₇ to C₁₈ substituted phenylalkylene, C₁ to C₁₂ heterocycloalkylene and C₁ to C₁₂ substituted heterocycloalkylene, -NH- and the formula:



wherein R⁹ and R¹⁰ are, independently, selected from the group consisting of a hydrogen atom, C₁ to C₁₂ alkyl, C₂ to C₁₂ alkenyl, C₂ to C₁₂ alkynyl, C₁ to C₁₂ substituted alkyl, C₂ to C₁₂ substituted alkenyl, C₂ to C₁₂ substituted alkynyl, C₁ to C₁₂ acyl, C₁ to C₁₂ substituted acyl, C₃ to C₇ cycloalkyl, C₃ to C₇ substituted cycloalkyl, C₅ to C₇ cycloalkenyl, C₅ to C₇ substituted cycloalkenyl, a heterocyclic ring, substituted heterocyclic ring, heteroaryl, substituted heteroaryl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂

heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, C₇ to C₁₈ phenylalkoxy, C₇ to C₁₈ substituted phenylalkoxy, phenyl, substituted phenyl, naphthyl, substituted naphthyl, cyclic C₂ to C₇ alkylene, substituted cyclic C₂ to C₇ alkylene, cyclic C₂ to C₇ heteroalkylene, substituted cyclic C₂ to C₇ heteroalkylene, carboxy, protected carboxy, hydroxymethyl and protected hydroxymethyl; and m and n are, independently, 0, 1, 2, 3 or 4; and

R⁷ and R⁸ are, independently, selected from the group consisting of a functionalized resin, a hydrogen atom, C₁ to C₁₂ alkyl, C₁ to C₁₂ substituted alkyl, phenyl, substituted phenyl, heterocycle, substituted heterocycle, C₃ to C₇ cycloalkyl, C₃ to C₇ substituted cycloalkyl, C₅ to C₇ cycloalkenyl, C₅ to C₇ substituted cycloalkenyl, C₂ to C₁₂ alkenyl, C₂ to C₁₂ substituted alkenyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, [[and]] C₁ to C₁₂ substituted heterocycloalkyl, C₁ to C₁₂ acyl, C₁ to C₁₂ substituted acyl, phenylsulfonyl, substituted phenylsulfonyl, C₁ to C₁₀ alkylsulfonyl, C₁ to C₁₀ substituted alkylsulfonyl, C₁ to C₁₂ alkylaminocarbonyl, C₁ to C₁₂ substituted alkylaminocarbonyl, phenylaminocarbonyl, substituted phenylaminocarbonyl, C₁ to C₁₂ alkylaminothiocarbonyl, C₁ to C₁₂ substituted alkylaminothiocarbonyl, phenylaminothiocarbonyl and substituted phenylaminothiocarbonyl;

provided that, where R⁶ is methylene, at least one of R¹ to R⁴ must be the formula -C(O)NR¹¹R¹²; or

provided that, where R⁶ is methylene, at least one of R¹ to R⁴ must be the formula -C(O)R¹¹, wherein R¹¹ is a heterocyclic ring or substituted heterocyclic ring, wherein said ring contains at least one nitrogen atom and wherein said nitrogen atom is attached to the carbonyl carbon; or a pharmaceutically acceptable salt of a compound thereof;

with the proviso that when R⁷ and R⁸ are hydrogen or -CH₂CH₃, substituents R¹, R², R³ and R⁴ cannot be hydrogen.

Claim 57 (new): The single compound of claim 56, wherein R¹, R² and R³ are, independently, selected from the group consisting of a hydrogen atom, halo, C₁ to C₁₂ alkyl, C₁

to C₁₂ substituted alkyl, carboxy, and the group consisting of (i) the formula $-C(O)NR^{11}R^{12}$ and (ii) the formula $-C(O)R^{11}$, wherein R¹¹ and R¹² are, independently, selected from the group consisting of a hydrogen atom, C₁ to C₁₂ alkyl, C₁ to C₁₂ substituted alkyl, C₂ to C₁₂ alkenyl, C₂ to C₁₂ substituted alkenyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, heterocycle and substituted heterocycle.

Claim 58 (new): The single compound of claim 56, wherein R¹, R², and R⁴ are each a hydrogen atom and R³ is selected from the group consisting of halo, C₁ to C₁₂ alkyl, C₁ to C₁₂ substituted alkyl, carboxy, and the group consisting of (i) the formula $-C(O)NR^{11}R^{12}$ and (ii) the formula $-C(O)R^{11}$, wherein R¹¹ and R¹² are, independently, selected from the group consisting of a hydrogen atom, C₁ to C₁₂ alkyl, C₁ to C₁₂ substituted alkyl, C₂ to C₁₂ alkenyl, C₂ to C₁₂ substituted alkenyl, C₇ to C₁₈ phenylalkyl, C₇ to C₁₈ substituted phenylalkyl, C₁ to C₁₂ heterocycloalkyl, C₁ to C₁₂ substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, heterocycle and substituted heterocycle.


Claim 59 (new): The single compound of claim 56, wherein R⁶ is the formula:

-D-W-E-

wherein:

W is absent or selected from the group consisting of phenylene, substituted phenylene, C₃ to C₇ cycloalkylene and C₃ to C₇ substituted cycloalkylene; and

D, which is directly attached to the nitrogen depicted in the formula, and E, which can be absent, are, independently, selected from the group consisting of C₁ to C₁₂ alkylene, C₁ to C₁₂ substituted alkylene, -NH- and the formula:



wherein R^9 and R^{10} are, independently, selected from the group consisting of a hydrogen atom, C_1 to C_{12} alkyl, C_1 to C_{12} substituted alkyl, C_3 to C_7 cycloalkyl, C_3 to C_7 substituted cycloalkyl, C_7 to C_{18} phenylalkyl, C_7 to C_{18} substituted phenylalkyl, phenyl, substituted phenyl; and m and n are, independently, 0, 1 or 2.

Claim 60 (new): The single compound of claim 56, wherein R^7 and R^8 are each a hydrogen atom.

Claim 61 (new): The single compound of claim 56, wherein:

R^1 , R^2 and R^4 are each a hydrogen atom and R^3 is the formula $-C(O)NR^{11}R^{12}$, wherein R^{11} is selected from the group consisting of a hydrogen atom, methyl, ethyl and benzyl and R^{12} is selected from the group consisting of a hydrogen atom, benzyl, 4-methoxyphenyl, 4-phenoxyphenyl, (1-ethyl-2-pyrrolidino)methyl, pyridin-2-ylmethyl, 2-(pyridin-2-yl)ethyl, methyl, 3,3,5-trimethylcyclohexyl, cyclohexyl, 3-(trifluoromethyl)benzyl, 6-indazolyl, 2-(ethoxycarbonyl)ethyl, ethoxycarbonylmethyl, cyclooctyl, cyclopropyl, (N,N-diethylamino)ethyl, 3-(2-oxo-1-pyrrolidino)propyl, (1-ethyl-2-pyrrolidinyl)methyl, pyridin-4-ylmethyl, 3-(4-morpholino)propyl, 4-methylphenyl, butyl and 2-thiazolyl;

R^5 is selected from the group consisting of 3-phenoxyphenyl, 3-hydroxy-4-methoxyphenyl, 4-acetamidophenyl, 4-phenoxyphenyl, 4-bromo-2-thienyl, 4-pyridyl, 2-butyl, 4-chloro-3-nitrophenyl, 3-nitrophenyl, 2,3-dichlorophenyl, 2,5-difluorophenyl, 5-methyl-2-furyl, 4-chloro-3-fluorophenyl, 2-phenyl-4-imidazolyl, 5-nitro-2-furyl, 4-bromophenyl, 2-norbornen-5-yl, 6-nitropiperonyl, 2-chloro-5-nitrophenyl, 5-hydroxy-2-nitrophenyl, 3-hydroxyphenyl, 3,4-difluorophenyl, 4-dimethylaminophenyl, 2-thienyl, 4-cyanophenyl, 3-cyanophenyl, 4-nitrophenyl, 2-fluorophenyl, 4-carboxyphenyl,

2-bromophenyl, 2-chloro-3,4-dimethoxyphenyl, 3-thienyl, 4-quinolyl, 4-methyl-5-imidazolyl, 4-hydroxyphenyl, 2-ethyl-5-formyl-4-methylimidazolyl, 4-chloro-2-nitrophenyl, 3-pyridyl, 3,4-dimethyl-6-nitrophenyl, 5-chloro-2-nitrophenyl and 2-nitrophenyl;

R⁶ is selected from the group consisting of methylenemethylene, ethylene, propylene, pentylene, isobutylenemethylene, 3-aminocarbonylpropylenemethylene, 2-methylthioethylenemethylene, isopropylenemethylene, phenylenemethylene, benzylenemethylene, cyclohexylenemethylene, 4-chlorobenzylenemethylene, indol-3-ylmethylene, 4-trifluoroacetamidobutylenemethylene, 3-guanidopropylenemethylene, -CH₂CH₂NH- and 1-cyclohexylene-4-NH-; and

R⁷ and R⁸ are each a hydrogen atom.

Claim 62 (new): The single compound of claim 56, wherein:

R¹, R² and R⁴ are each a hydrogen atom and R³ is the formula -C(O)R¹¹, wherein R¹¹ is selected from the group consisting of 1,3,3-trimethyl-6-aza-6-bicyclo(3,2,1)octyl, 4-(4-fluorophenyl)-1-piperazino, 4-acetyl-1-piperazino, morpholino, 2-methyl-4-(3-methylphenyl)-1-piperazino, 4-ethoxycarbonylpiperidino and N-methylhomopiperazino;

R⁵ is selected from the group consisting of 3-phenoxyphenyl, 3-hydroxy-4-methoxyphenyl, 4-acetamidophenyl, 4-phenoxyphenyl, 4-bromo-2-thienyl, 4-pyridyl, 2-butyl, 4-chloro-3-nitrophenyl, 3-nitrophenyl, 2,3-dichlorophenyl, 2,5-difluorophenyl, 5-methyl-2-furyl, 4-chloro-3-fluorophenyl, 2-phenyl-4-imidazolyl, 5-nitro-2-furyl, 4-bromophenyl, 2-norbornen-5-yl, 6-nitropiperonyl, 2-chloro-5-nitrophenyl, 5-hydroxy-2-nitrophenyl, 3-hydroxyphenyl, 3,4-difluorophenyl, 4-dimethylaminophenyl, 2-thienyl, 4-cyanophenyl, 3-cyanophenyl, 4-nitrophenyl, 2-fluorophenyl, 4-carboxyphenyl, 2-bromophenyl, 2-chloro-3,4-dimethoxyphenyl, 3-thienyl, 4-quinolyl, 4-methyl-5-imidazolyl, 4-hydroxyphenyl, 2-ethyl-5-formyl-4-methylimidazolyl, 4-chloro-2-nitrophenyl, 3-pyridyl, 3,4-dimethyl-6-nitrophenyl, 5-chloro-2-nitrophenyl and 2-nitrophenyl;

R⁶ is selected from the group consisting of methylenemethylene, ethylene, propylene, pentylene, isobutylenemethylene, 3-aminocarbonylpropylenemethylene, 2-methylthioethylenemethylene,

isopropylmethylene, phenylmethylene, benzylmethylene, cyclohexylmethylmethylene, 4-chlorobenzylmethylene, indol-3-ylmethylmethylene, 4-trifluoroacetamidobutylmethylene, 3-guanidopropylmethylene, $-\text{CH}_2\text{CH}_2\text{NH}-$ and 1-cyclohexylene-4-NH-; and

R^7 and R^8 are each a hydrogen atom.

Claim 63 (new): The single compound of claim 56, wherein:

R^1 , R^2 and R^4 are each a hydrogen atom and R^3 is the formula $-\text{C}(\text{O})\text{NR}^{11}\text{R}^{12}$, wherein R^{11} is selected from the group consisting of a hydrogen atom, methyl, ethyl and benzyl and R^{12} is selected from the group consisting of a hydrogen atom, 2-(2-methoxyphenyl)ethyl, (1-ethyl-2-pyrrolidino)methyl, pyridin-2-ylmethyl, 2-methyl-5-chlorophenyl, (2-(pyridin-2-yl)ethyl), 1-ethyl-2-pyrrolidinylmethyl, 3,3,5-trimethylcyclohexyl, 3,4-methylenedioxyphenyl, 3-(trifluoromethyl)benzyl, pyridin-4-ylmethyl, 6-indazolyl, 2-(ethoxycarbonyl)ethyl, cyclooctyl, cyclopropyl, benzyl, N,N-(diethylamino)ethyl, 3-(2-oxo-1-pyrrolidine)propyl, 3-(4-morpholino)propyl, (ethoxycarbonyl)methyl and cyclohexyl;

R^5 is selected from the group consisting of phenoxyphenyl, 4-hydroxy-3-methoxyphenyl, 3,4,5-trimethoxyphenyl, 3-hydroxy-4-methoxyphenyl, 4-acetamidophenyl, 4-phenoxyphenyl, 4-methoxyl-1-naphthyl, 4-bromo-2-thienyl, 4-pyridyl, isopropyl, 2-methylthioethyl, 4-chloro-3-nitrophenyl, 3-nitrophenyl, 4-t-butylphenyl, 2,3-dichlorophenyl, 3,5-bis(trifluoromethyl)phenyl, 2,5-difluorophenyl, 2-quinolyl, 2-chloro-3,4-dimethoxyphenyl, 5-methyl-2-furyl, 4-chloro-3-fluorophenyl, 2-phenyl-4-imidazolyl, 2-(ethoxycarbonyl)cyclopropyl, 5-nitro-2-furyl, 4-bromophenyl, cyclopropyl, 2-norbornen-5-yl, 6-nitropiperonyl, 2-chloro-5-nitrophenyl, 5-hydroxy-2-nitrophenyl, 3-hydroxyphenyl, 3,4-difluorophenyl, 4-dimethylaminophenyl, 4-methylthiophenyl, 4-(trifluoromethyl)phenyl, 2-thienyl, 2,3-dimethoxyphenyl, 3-ethoxy-4-hydroxyphenyl, 4-cyanophenyl, 3-cyanophenyl, 2-furyl, 4-nitrophenyl, 1-naphthyl, 2-methoxyphenyl, 4-isopropylphenyl, piperonyl, 2-fluorophenyl, 4-ethoxyphenyl and 2,4-dihydroxyphenyl;

R⁶ is selected from the group consisting of methylene, ethylidene, ethylene, propylene, pentylene, isopentylidene, 3-aminocarbonylbutylidene, 2-methylthiopropylidene, isobutylidene, phenylmethylene, benzylmethylene, cyclohexylethylidene, 4-chlorobenzylmethylene, indol-3-ylethylidene, 4-trifluoroacetamidopentylidene, 3-guanidobutylidene, hydroxyethylidene, 2-aminocarbonylpropylidene, isopentylidene, mercaptoethylidene, 4-hydroxybenzylmethylene, 1,3-phenylene, 1,4-phenylene, 1,4-(phenylene)-NH-, 3,6-dioxaoctylene-NH-, -CH₂CH₂NH- and 1,4-(cyclohexylene)-NH-and

R⁷ and R⁸ are each a hydrogen atom.

Claim 63 (new): The single compound of claim 56, wherein:

R¹, R² and R⁴ are each a hydrogen atom and R³ is the formula -C(O)R¹¹, wherein R¹¹ is selected from the group consisting of 1,3,3-trimethyl-6-aza-6-bicyclo(3,2,1)octyl, 4-(4-fluorophenyl)-1-piperazino, 4-acetyl-1-piperazino, piperazino, 2-methyl-4-(3-methylphenyl)-1-piperazino, 4-(ethoxycarbonyl)piperidino, N-methylhomopiperazino and N,N'-diisopropylimidamino;

R⁵ is selected from the group consisting of phenoxyphenyl, 4-hydroxy-3-methoxyphenyl, 3,4,5-trimethoxyphenyl, 3-hydroxy-4-methoxyphenyl, 4-acetamidophenyl, 4-phenoxyphenyl, 4-methoxyl-1-naphthyl, 4-bromo-2-thienyl, 4-pyridyl, isopropyl, 2-methylthioethyl, 4-chloro-3-nitrophenyl, 3-nitrophenyl, 4-t-butylphenyl, 2,3-dichlorophenyl, 3,5-bis(trifluoromethyl)phenyl, 2,5-difluorophenyl, 2-quinolyl, 2-chloro-3,4-dimethoxyphenyl, 5-methyl-2-furyl, 4-chloro-3-fluorophenyl, 2-phenyl-4-imidazolyl, 2-(ethoxycarbonyl)cyclopropyl, 5-nitro-2-furyl, 4-bromophenyl, cyclopropyl, 2-norbornen-5-yl, 6-nitropiperonyl, 2-chloro-5-nitrophenyl, 5-hydroxy-2-nitrophenyl, 3-hydroxyphenyl, 3,4-difluorophenyl, 4-dimethylaminophenyl, 4-methylthiophenyl, 4-(trifluoromethyl)phenyl, 2-thienyl, 2,3-dimethoxyphenyl, 3-ethoxy-4-hydroxyphenyl, 4-cyanophenyl, 3-cyanophenyl, 2-furyl, 4-nitrophenyl, 1-naphthyl, 2-methoxyphenyl, 4-isopropylphenyl, piperonyl, 2-fluorophenyl, 4-ethoxyphenyl and 2,4-dihydroxyphenyl;

R^6 is selected from the group consisting of methylene, ethylidene, ethylene, propylene, pentylene, isopentylidene, 3-aminocarbonylbutylidene, 2-methylthiopropylidene, isobutylidene, phenylmethylene, benzylmethylene, cyclohexylethylidene, 4-chlorobenzylmethylene, indol-3-ylethylidene, 4-trifluoroacetamidopentylidene, 3-guanidobutylidene, hydroxyethylidene, 2-aminocarbonylpropylidene, isopentylidene, mercaptoethylidene, 4-hydroxybenzylmethylene, 1,3-phenylene, 1,4-phenylene, 1,4-(phenylene)-NH-, 3,6-dioxaoctylene-NH-, $-CH_2CH_2NH-$ and 1,4-(cyclohexylene)-NH-; and

R^7 and R^8 are each a hydrogen atom.

Claim 64 (new): The single compound of claim 56, wherein

R^1 , R^2 , R^4 , R^7 and R^8 are each a hydrogen atom;

R^3 is the formula $-C(O)NR^{11}R^{12}$, wherein R^{11} is a hydrogen atom and R^{12} is selected from the group consisting of pyridin-2-ylmethyl and 3,3,5-trimethylcyclohexyl;

R^5 is selected from the group consisting of 4-N,N-dimethylaminophenyl, 5-chloro-2-nitrophenyl, 4-bromo-2-thienyl, 2-butyl, 5-nitro-2-furyl, 4-bromophenyl, 2-thienyl, 3-thienyl, 3-cyanophenyl, 4-cyanophenyl, 4-quinolyl and 4-hydroxyphenyl; and

R^6 is methylene.

Claim 65 (new): The single compound of claim 56, wherein R^4 is selected from the group consisting of a hydrogen atom, halo, hydroxy, protected hydroxy, cyano, C_1 to C_{12} alkyl, C_2 to C_{12} alkenyl, C_2 to C_{12} alkynyl, C_1 to C_{12} substituted alkyl, C_2 to C_{12} substituted alkenyl, C_2 to C_{12} substituted alkynyl, C_1 to C_{12} alkoxy, C_1 to C_{12} substituted alkoxy, C_1 to C_{12} acyloxy, C_1 to C_{12} acyl, C_3 to C_7 cycloalkyl, C_3 to C_7 substituted cycloalkyl, C_5 to C_7 cycloalkenyl, C_5 to C_7 substituted cycloalkenyl, heterocyclic ring, substituted heterocyclic ring, C_7 to C_{18} phenylalkyl, C_7 to C_{18} substituted phenylalkyl, C_1 to C_{12} heterocycloalkyl, C_1 to C_{12} substituted heterocycloalkyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, cyclic C_2 to C_7 alkylene, substituted cyclic C_2 to C_7 alkylene, cyclic C_2 to C_7 heteroalkylene, substituted cyclic C_2 to C_7 heteroalkylene.

Claim 66 (new): The single compound of claim 56, wherein R^5 is selected from the group consisting of phenyl, substituted phenyl, C_1 to C_{12} heterocycloalkyl, C_1 to C_{12} substituted heterocycloalkyl, heterocycle, substituted heterocycle, naphthyl, substituted naphthyl, C_3 to C_7 cycloalkyl, C_3 to C_7 substituted cycloalkyl, C_5 to C_7 cycloalkenyl and C_5 to C_7 substituted cycloalkenyl.

Claim 67 (new): The single compound of claim 42, wherein R^6 is methylene and at least one of R^1 to R^4 must be the formula $-C(O)NR^{11}R^{12}$.

Claim 68 (new): The single compound of claim 42, wherein R^6 is methylene and at least one of R^1 to R^4 must be the formula $-C(O)R^{11}$, where R^{11} is a heterocyclic ring or substituted heterocyclic ring, wherein said ring contains at least one nitrogen atom and where said nitrogen atom is attached to the carbonyl carbon.

Claim 69 (new): The single compound of claim 44, wherein R^6 is methylene and at least one of R^1 to R^4 must be the formula $-C(O)NR^{11}R^{12}$;

Claim 70 (new): The single compound of claim 44, wherein R^6 is methylene and at least one of R^1 to R^4 must be the formula $-C(O)R^{11}$, wherein R^{11} is a heterocyclic ring or substituted heterocyclic ring, wherein said ring contains at least one nitrogen atom and wherein said nitrogen atom is attached to the carbonyl carbon.